

15th January 2009

Is the Unitarity Triangle Right?

P. F. Harrison¹ and D. R. J. Roythorne
Department of Physics, University of Warwick,
Coventry CV4 7AL. UK

and

W. G. Scott
Rutherford Appleton Laboratory,
Chilton, Didcot, Oxon OX11 0QX. UK

Abstract

The latest fits to the CKM matrix indicate that $\alpha = (90.7^{+4.5}_{-2.9})^\circ$. The proximity of α to a right-angle raises the question: is it merely accidental or is it due to some physics beyond the Standard Model? In the framework of our recently-proposed flavour permutation symmetry, we consider the similarities between the quark and lepton mixing matrices, V and U , arguing that the relative smallness of one element in each suggests common constraints. These constraints link the smallness of V_{ub} and U_{e3} with each other, and with the approximate $\mu - \tau$ symmetry observed in leptonic mixing, together with a prediction of a large Dirac CP phase in both the quark and lepton sectors. In the quark case, we predict $\alpha = (89.0 \pm 0.2)^\circ$, in agreement with data and suggesting that the unitarity triangle is in fact very nearly, but not exactly right.

Talk given at the International Conference on Particles and Nuclei (PANIC08), Eilat, Israel, 13th November 2008.²

¹Speaker and Corresponding Author, E-mail: p.f.harrison@warwick.ac.uk

²<http://www.weizmann.ac.il/MaKaC/contributionDisplay.py?contribId=235&sessionId=41&materialId=slides&confId=0>

I have to thank Bjorken for the title of my talk, since he posed just this question in 1997 [1]. He was interested, together with Stech [2], in the possibility that the angle γ of the unitarity triangle (UT) might be exactly 90° . He also commented (negatively) on an earlier prediction by Fritzsche and Xing [3] that $\alpha \sim 90^\circ$ might be preferred. The latest fits [4] to the CKM matrix using data from the B factories and other facilities indicate that $\alpha = (90.7^{+4.5}_{-2.9})^\circ$, so that Fritzsche and Xing may indeed have been correct. Their successful prediction was based on their “four texture zero” model of hermitian quark mass matrices which needed α in the range $\sim 90^\circ \pm 20^\circ$ to be consistent with the famous relationship: $\sin^2 \theta_C \simeq \frac{m_d}{m_s} + \frac{m_u}{m_c}$. We proposed in 2007 [5] an alternative explanation for $\alpha \simeq 90^\circ$.

We first introduced the idea [5] of flavour-symmetric mixing observables (FSMOs), namely those mixing observables having specific transformation properties (either even or odd) under the discrete group of (separate) permutations of the rows and columns of the mixing matrix (for either quarks or leptons). FSMOs share this flavour-symmetry property with the Jarlskogian [6], J , which is the prototype flavour-odd observable (with J^2 a prototype flavour-even variable). FSMOs were also shown to be expressible in terms of simple functions of the relevant mass matrices. We now propose [7], a complete set of such variables based on the K -matrix [8] of mixing observables, $K_{\alpha i} := \text{Re}(V_{\beta j} V_{\gamma k} V_{\beta k}^* V_{\gamma j}^*)$, which is the CP -conserving analogue of J :

$$\begin{aligned} S &:= \sum_{\gamma k} K_{\gamma k}; & \mathcal{R}' &:= \sum_{\gamma k} (K_{\alpha i} K_{\beta j} + K_{\alpha j} K_{\beta i}); \\ \mathcal{K} &:= \text{Det } K; & 3J^2 &= \sum_{\alpha} \sum_{i \neq j} K_{\alpha i} K_{\alpha j} = \sum_i \sum_{\alpha \neq \beta} K_{\alpha i} K_{\beta i}. \end{aligned} \quad (1)$$

S , $\mathcal{R}'$ and J^2 are flavour-even, while $\mathcal{K}$ is odd. These four variables completely specify the physical content of complex 3×3 mixing matrices, modulo permutations of rows and columns and the sign of J . Expressions for them in terms of the standard (PDG) mixing parameters, the Wolfenstein parameters and our P -matrix based FSMOs [5] are given in [7]. We note the following suggestive hierarchy among them: i) CP -phase $\delta = 0 \Rightarrow J = 0$; ii) Any one mixing angle, $\theta_{ij} = 0 \Rightarrow J = \mathcal{K} = 0$; iii) Any two $\theta_{ij} = 0 \Rightarrow J = \mathcal{K} = \mathcal{R}' = 0$; iv) All three $\theta_{ij} = 0 \Rightarrow J = \mathcal{K} = \mathcal{R}' = S = 0$.

It is often emphasised that the quark and lepton mixing matrices have very different forms. However, a unified description remains an important goal, which we seek in flavour-symmetric terms [5]. Both the quark and lepton mixing matrices share the common feature that their top right-hand element is significantly smaller than the typical magnitudes of their other elements. Hence we seek a flavour-symmetric condition for one small mixing matrix element. Now, a mixing matrix has a zero element if and only if:

$$\mathcal{K} = 0 \quad \text{and} \quad J = 0. \quad (2)$$

We recall that with μ - τ -reflection symmetry [9] (which holds at least to a good approximation in the leptonic case), the constraint $\mathcal{K} = 0$ follows trivially from the elementary properties of determinants, since in that case, the μ and τ rows of the K matrix are equal to each other. Moreover, both conditions, Eq. (2), hold exactly in tri-bimaximal mixing [10]. The two conditions, Eq. (2), may be combined, both being implied when the product, $\mathcal{P}$, of

the moduli-squared of all nine elements of the mixing matrix satisfies:

$$\mathcal{P} = 0, \quad \text{where} \quad \mathcal{P} \equiv \prod_{\alpha i} |V_{\alpha i}|^2 = \mathcal{K}^2 + J^2(2J^2 + \mathcal{K}')^2, \quad (3)$$

which is zero iff $\mathcal{K} = 0$ and $J = 0$ (since $\mathcal{K}' \geq 0$ always).

For a small, rather than a zero mixing matrix element, one or other (or both) of the conditions, Eq. (2), should be relaxed. Further, since for the quarks, clearly $J \neq 0$, while for both quarks and leptons, the data are consistent with $\mathcal{K} = 0$, we are led to the following predictive conjecture, consistent with both the quark and lepton data:

$$\mathcal{K} = 0 \quad \text{and} \quad |J/J_{\max}| = \text{small} \quad (4)$$

(it is not implied that the small quantity necessarily has the same value in both sectors). Equation (4) is a unified and flavour-symmetric, partial description (in the sense that only two degrees of freedom are constrained) of both lepton and quark mixing matrices. It implies the existence of at least one small element in each mixing matrix, as manifested by U_{e3} and V_{ub} , and is associated with $\mu - \tau$ symmetry [9], which is suggested by the lepton data.

While relaxing slightly *both* constraints, Eq. (2), allows a small mixing angle, θ (eg. θ_{13}), and an arbitrary CP -phase, relaxing instead *only* the J -constraint while keeping $\mathcal{K} = 0$, Eq. (4), still yields a small mixing angle, θ , but now subject to the phase-condition that one UT angle $\rightarrow 90^\circ$, as $J \rightarrow 0$ [such that $(\theta/\theta_{ij}) \rightarrow 0$ (for $\theta_{ij} \neq \theta$)]. Moreover, since the $(\theta_{13}/\theta_{ij})$ are small for both the quarks and leptons, the deviations from 90° are small and calculable in terms of them. Flavour symmetry prevents an a priori prediction of *which* UT angle is $\simeq 90^\circ$, but this can be obtained from the data (in general, flavour symmetry is spontaneously broken by the solutions of flavour-symmetric constraint equations).

For the quarks, we use our formula [7, 11] for $\mathcal{K}$ in the Wolfenstein parameterisation (being based on data, the latter already breaks flavour symmetry) to show that $\mathcal{K} = 0 \Rightarrow \alpha \simeq 90^\circ$. Defining the squares of the two non-trivial sides of the standard B physics unitarity triangle as $u \equiv \bar{\rho}^2 + \bar{\eta}^2$ and $v \equiv 1 - 2\bar{\rho} + \bar{\rho}^2 + \bar{\eta}^2$, the cosine rule gives $\cos \alpha = \frac{u+v-1}{2\sqrt{uv}}$. Meanwhile $\mathcal{K} = 0$ implies [7, 11]:

$$u + v - 1 \equiv 2(\bar{\rho}^2 + \bar{\eta}^2 - \bar{\rho}) = 2\lambda^2 uv \Rightarrow \cos \alpha = \lambda^2 \sqrt{uv} \simeq \lambda^2 \bar{\eta} \quad (5)$$

at leading order in λ^2 (it vanishes as $\frac{\sin \theta_{13}}{\sin \theta_{23}} \sim \lambda \sqrt{u} \rightarrow 0$, as foreseen). Eq. (5) predicts:

$$(90^\circ - \alpha) \simeq \lambda^2 \bar{\eta} = 1.0^\circ \pm 0.2^\circ, \quad (6)$$

compared with its current measured value [4]: $(90^\circ - \alpha) = 2_{-6}^{+5}^\circ$ (direct measurement) or $(90^\circ - \alpha) = -1_{-5}^{+3}^\circ$ (full CKM fit). The unitarity triangle is indeed, very nearly right! It will be interesting to test Eq. (6) more precisely in future B physics experiments, such as LHCb and super flavour factories.

We know experimentally that for leptons, U_{e3} is the only small mixing matrix element. Hence, only the UT angles $\phi_{\mu 1}$, $\phi_{\tau 1}$, $\phi_{\mu 2}$ and $\phi_{\tau 2}$ (using the nomenclature of [11, 12]) can be close to 90° , and they all satisfy (see eg. Fig. 1 of [12]) $|\cos \phi_{\alpha i}| \simeq |\cos \delta|$, where

$\delta = \text{Arg}(U_{e3})$. Using our formula [7] for $\mathcal{K}$ in terms of PDG parameters, we find that $\mathcal{K} = 0$ implies:

$$|90^\circ - \delta| \simeq 4 \cot 2\theta_{12} \cot 2\theta_{23} \sin \theta_{13} \simeq 2\sqrt{2} \sin \theta_{13} \sin(\theta_{23} - 45^\circ) \lesssim 4^\circ, \quad (7)$$

(1σ) at leading order in small quantities (again vanishing as $\frac{\sin \theta_{13}}{\sin \theta_{23}} \rightarrow 0$ as expected). Our conjecture thus predicts a large CP -violating phase in the MNS matrix, which is promising for the discovery of leptonic CP violation at eg. a future Neutrino Factory.

References

- [1] J.D. Bjorken, Summary talk at the 2nd International Conference on B Physics and CP Violation, Hawaii (1997), hep-ph/9706524.
- [2] B. Stech, Phys. Rev. Lett. B130 (1983) 189.
- [3] H. Fritzsch and Z.Z. Xing, Phys. Lett. B353 (1995) 114, hep-ph/9502297.
- [4] CKMfitter Group (J. Charles et al.), Eur. Phys. J. C41 (2005) 1, hep-ph/0406184, updated results and plots: <http://ckmfitter.in2p3.fr>; M. Bona et al. (UTfit Collaboration), hep-ph/0701204.
- [5] P.F. Harrison, D.R.J. Roythorne and W.G. Scott, Phys. Lett. B657 (2007) 210, arXiv:0709.1439.
- [6] C. Jarlskog, Phys. Rev. Lett. 55 (1985) 1039; Z. Phys. C29 (1985) 491.
- [7] P.F. Harrison, D.R.J. Roythorne and W.G. Scott, in preparation.
- [8] P.F. Harrison, W.G. Scott and T.J. Weiler, Phys. Lett. B641 (2006) 372, hep-ph/0607335.
- [9] C.S. Lam, Phys. Lett. B507 (2001) 214, hep-ph/0104116; M. Kitabayashi and M. Yatsue, Phys. Lett. B524 (2002) 308, hep-ph/0110303; P.F. Harrison and W.G. Scott, Phys. Lett. B547 (2002) 219, hep-ph/0210197; W. Grimus and L. Lavoura, Phys. Lett. B579 (2004) 113.
- [10] P.F. Harrison, D.H. Perkins, and W.G. Scott, Phys. Lett. B458 (1999) 79, hep-ph/9904297; Phys. Lett. B530 (2002) 167, hep-ph/0202074.
- [11] P.F. Harrison, S. Dailison and W.G. Scott, arXiv:0904.3077.
- [12] J.D. Bjorken, P.F. Harrison and W.G. Scott, Phys. Rev. D **74** (2006) 073012 hep-ph/0511201.